arXiv:1212.1032v1 [cond-mat.stat-mech] 5 Dec 2012

Correlations in the $n \to 0$ limit of the dense $O(n)$ loop model

V.S. Poghosyan[1] and V.B. Priezzhev[2]

[1]*Institute for Informatics and Automation Problems*
NAS of Armenia, 0014 Yerevan, Armenia
[2]*Bogoliubov Laboratory of Theoretical Physics,*
Joint Institute for Nuclear Research, 141980 Dubna, Russia

The two-dimensional dense $O(n)$ loop model for $n = 1$ is equivalent to the bond percolation and for $n = 0$ to the dense polymers or spanning trees. We consider the boundary correlations on the half space and calculate the probability P_b that a cluster of bonds has a single common point with the boundary. In the limit $n \to 0$, we find an analytical expression for P_b using the generalized Kirchhoff theorem. The obtained exact value is surprisingly close to the value of P_b conjectured for the dense $O(1)$ loop model.

Keywords: dense $O(n)$ loop model, bond percolation, spanning trees, correlation functions.

I. INTRODUCTION

The dense $O(n)$ loop model [1] is defined by drawing two lines in each elementary cell of the square lattice. Two possible states of the cell are shown in Fig.1. The lines on the whole lattice with appropriate boundary conditions form a system of closed loops with the Boltzmann weight n ascribed to every loop.



FIG. 1: Elementary cells.

A rise of interest to the $O(n)$ model, especially to the case $n = 1$, was triggered by remarkable observations concerning the ground state of the antiferromagnetic XXZ quantum



FIG. 2: Loops on the horizontally periodic lattice.

chain and, equivalently, to properties of the transfer matrix of dense loop model taken in the Hamiltonian limit of extreme spatial anisotropy [2–7]. It was noted that the elements of the ground state of the loop Hamiltonian are equal to the cardinality of different subsets of configurations of the Fully Packed Loop (FPL) model which in their turn are connected with classes of Alternating Sign Matrices (ASM).

The states of the $O(n)$ model can be defined in terms of the connectivity condition for points of intersection between loops and a horizontal line cutting the loops at these points. Two points are connected by a link if there exists a line between them via the half space above the cut. For instance, imposing periodic boundary conditions in horizontal direction for the lattice in Fig.2 and specifying the boundary conditions at the upper edge, we obtain three minimal links between points 1 and 6, 2 and 3, 4 and 5 which are points of intersection between loops and the bottom line of lattice belonging to upper half space. A typical configuration of links for a larger lattice is given in Fig.3.

Several interesting conjectures have been obtained about the probability distribution of configurations of links of the dense $O(1)$ loop model. To be closer to the aim of this article, we mention just one conjecture about the number of configurations containing the minimal



FIG. 3: A link configuration.

link at a given position [7]

$$\sum \ldots \frown \ldots = \frac{3}{8}\frac{N^2}{N^2-1}A_{HT}(N) \tag{1.1}$$

for even periodic system and

$$\sum \ldots \frown \ldots = \frac{3}{8}\frac{N^2-1}{N^2}A_{HT}(N) \tag{1.2}$$

for odd periodic system. Here $A_{HT}(N)$ is the number of $N \times N$ half turn invariant ASM [8].

Since the total number of periodic link configurations is $A_{HT}(N)$, the probability of minimal link $Prob(\ldots \frown \ldots)$ tends to 3/8 in the limit $N \to \infty$. Due to the simple bijection between the loop configurations on the square lattice and the bond configurations on its sublattices, the dense $O(1)$ model is equivalent to to the bond percolation. Then, the constant 3/8 has a simple interpretation as the probability of a certain class of percolation clusters (see Section II). Despite its simplicity, an analytical derivation of the probability of minimal links as well as of other link configurations remains an open problem.

In this work, we consider the problem of evaluation of $Prob(\ldots \frown \ldots)$ in the dense $O(n)$ model for the case $n = 0$. In the limit $n \to 0$, the bulk loops disappear and the system of lines is converted into the model of dense polymers [9]. The bijection giving the bond percolation for $n = 1$ on a sublattice, leads in the limit $n \to 0$ to the spanning trees on the same sublattice. A lot is known about correlation functions of the spanning tree model due to the Kirchhoff theorem [10]. However, we will see in subsequent sections that evaluation of $Prob(\ldots \frown \ldots)$ involves non-local diagrams which need a generalization of the Kirchhoff determinant formula. Similar correlations appear in the Abelian sandpile theory [11–15], in the problem of loop-erazed random walk [16, 17], in the problem of "watermelon" embedded into the spanning tree [18, 19]. Below, we use the generalized determinant formula to find $Prob(\ldots \frown \ldots)$ and compare the result obtained for $n = 0$ with that conjectured for $n = 1$.

II. LINK STRUCTURE AND BOUNDARY CLUSTERS

The square lattice of sites with integer coordinates (n, m) can be divided into two sublattices, black and white. For sites of the black sublattice $n + m$ is even, for sites of the white one $n + m$ is odd. The bijection between loop and bond configurations mentioned in Introduction is shown in Fig.4. The neighboring sites of each sublattice are connected by a bond if it does not intersect the lines of elementary cell. Each connected cluster of bonds in the bulk of lattice is situated inside a loop. Each bulk cluster on the black sublattice is surrounded by a connected cluster of bonds on the white sublattice and vice versa.



FIG. 4: The bijection between bonds and elementary cells.

The horizontal line cutting the loops cuts bond clusters inside them and produces boundary clusters surrounded by pieces of loops. Every piece has two points of intersections with the horizontal line and corresponds to the link between these points. The clusters of bonds corresponding to the loop configuration in Fig.2 are shown in Fig.5.

Since we are interested in the limiting case of large lattice $N \to \infty$, we take the infinite horizontal line and neglect the left and right boundary conditions. We put the origin of the lattice at a selected black site on the horizontal line and consider the boundary clusters of bonds containing the origin. The boundary clusters can be classified into four types shown schematically in Fig.6. The cluster of type (a) consists of a single vertex coinciding with the origin. The type (b) represents clusters containing no points of the horizontal line besides the origin. The clusters of type (c) contain the origin and one or more boundary vertices left or right from the origin. The clusters of type (d) contain the origin and an arbitrary number of boundary vertices left and right from the origin.

The minimal link probability $Prob(\ldots \frown \ldots)$ is the sum of probabilities P_a of the clusters of type (a) and P_b of the clusters of type (b) because the minimal link corresponds to the piece of loop surrounding the origin $(0, 0)$ together with the cluster of bonds attached to the



FIG. 5: The bond configuration corresponding to the loop configuration from Fig.2.



FIG. 6: Four types of the boundary clusters.

origin.

In the $O(1)$ model, the probability P_a is elementary because the isolated site $(0, 0)$ of the black sublattice corresponds to two bonds on the dual white sublattice: one bond connecting sites $(-1,0)$ and $(0,1)$ and another one connecting sites $(0,1)$ and $(1,0)$. For the bond percolation, where the bond probability is $1/2$, $P_a = 1/4$. The conjectured value of $Prob(\ldots \frown \ldots)$ is $3/8$, then $P_b = 1/8$. Considering the system of links as a height profile [6], one can note that the minimal links correspond to peaks of the profile and the local link configurations produced by clusters of type (d) correspond to valleys of the profile. The numbers of peaks and valleys in the periodic system coincide, so we have $P_d = 3/8$. Using the identity

$$P_a + P_b + 2P_c + P_d = 1 \tag{2.1}$$

we conclude that $P_c = 1/8$. The symmetry $P_b = P_c$ is wonderful because the bond clusters of type (b),(c) do not obey any visible symmetry in the percolation theory. In Table 1, we show the results of enumeration of clusters of type (a),(b),(c) and (d) for the strip of increasing size. We see from Fig.7 that the convergence of $P_a + P_b$ to $3/8$ is of an order of $1/N^2$ as it is expected from the conjecture (1.1).



FIG. 7: Monte-Carlo simulation result. Number of samples is 10^8, the $N \times N$ upper-left half-plane with $N = 3, 5, 7, \ldots, 41$ is considered.

Our aim in this work is calculation of probabilities $P_a,\ P_b,\ P_c,\ P_d$ for the free-fermion $O(0)$ model. Elimination of loops in the limit $n \to 0$ converts the wavy lines of the $O(n)$ model into dense polymers [9] and the bond clusters on the black and white sublattices into spanning trees on these sublattices. A spanning tree on the black (white) sublattice is a

N	P_d	$P_a + P_b$	P_c
3	0.0937471400	0.5313127300	0.1874740000
5	0.1675092500	0.4423504000	0.1950355800
7	0.2124676800	0.4119452400	0.1877856200
9	0.2418490300	0.3982314800	0.1799565400
11	0.2623780800	0.3909652700	0.1733454800
13	0.2774548300	0.3866752600	0.1679902400
15	0.2890090100	0.3838207700	0.1636097200
17	0.2981210300	0.3818828100	0.1599813100
19	0.3055246300	0.3805543900	0.1569469100
21	0.3114667000	0.3797553600	0.1542892800
23	0.3166525700	0.3788883400	0.1522516400
25	0.3211872800	0.3781053000	0.1503894000
27	0.3247541400	0.3778676500	0.1486765700
29	0.3281502900	0.3774419200	0.1472263800
31	0.3308674800	0.3771801800	0.1460154300
33	0.3334175200	0.3769390100	0.1447843200
35	0.3357127000	0.3766440200	0.1438112700
37	0.3377116700	0.3765149400	0.1428866100
39	0.3395580300	0.3764257200	0.1420234400
41	0.3412365600	0.3762179600	0.1413121400
∞	$0.37497 \approx 3/8$	$0.37499 \approx 3/8$	$0.12506 \approx 1/8$

TABLE I: The dense $O(1)$ loop model: values for $N = \infty$ are obtained from the extrapolation (see Fig.7).

connected cluster containing all black (white) sites. Due to the cut, the connected spanning tree splits into separated components attached to the horizontal line. The classification of the components (Fig.6) is the same as in the percolation model. For definiteness, we consider black components and calculate their statistics via probabilities of white components surrounding black clusters. Normally, one selects an arbitrary site of the tree as a root. Then all bonds of the tree become oriented towards the root. It is convenient to choose the root of tree on the white sublattice at a site below the horizontal line that cuts the lattice. Then a white cluster having k sites on the cut can be oriented by k different ways because every boundary site can be connected with the root by a path along the tree. Evaluation of P_a is almost as elementary as that for the $O(1)$ case. Indeed, we have to find the probabilities of three configurations of two oriented bonds on the dual white sublattice:

1a. The bonds from $(-1,0)$ to $(0,1)$ and from $(0,1)$ to $(1,0)$;

2a. The bonds from $(1,0)$ to $(0,1)$ and from $(0,1)$ to $(-1,0)$;

3a. The bonds from $(-1,0)$ to $(0,1)$ and from $(1,0)$ to $(0,1)$.

The probability P_a is the sum of these probabilities, $P_a = P_{1a} + P_{2a} + P_{3a}$.

Evaluation of P_b is a more delicate problem. The cluster of type (b) on the black sublattice is surrounded by bonds of a cluster on the white sublattice. The bonds are directed and their sequence gives continuous paths on the white sublattice of three different forms:

1b. The self-avoiding path of an arbitrary length which starts at the site $(-1,0)$, visits the site $(1,0)$, either stops at $(1,0)$ or continues throughout the white sublattice above the cut and ends at a white site on the cut.

2b. The path starts at the site $(1,0)$, visits the site $(-1,0)$, either stops at $(-1,0)$ or continues throughout the white sublattice above the cut and ends at a white site on the cut.

3b. Two self-avoiding paths start at the sites $(-1,0)$ and $(1,0)$, meet at a white site s_0 above the cut and then the single self-avoiding path continues from s_0 up to the end at a white site on the cut.

Denoting the probabilities of these situations by P_{1b}, P_{2b}, P_{3b}, we have to find $P_b = P_{1b} + P_{2b} + P_{3b} - P_a$.

While the probabilities constituting P_a are local correlation functions and can be calculated within a usual approach of the perturbed Laplacian, the probabilities in P_b are essentially non-local and their calculation needs a special technique. Before formulating these technical tools, we chose more convenient coordinates for our problem.

Both black and white sublattices are square lattices with the elementary cell $\sqrt{2} \times \sqrt{2}$ that are turned by $\pi/4$ with respect to the basic lattice. We chose the standard orientation for the sublattices and put the step of these square lattices to be equal 1. In new coordinates, the former cut is the diagonal line crossing the origin. The sites of black sublattice belonging to cut have coordinates $(0,0)$, $(1,1)$, $(-1,-1)$, $(2,2)$, $(-2,-2), \ldots$. The coordinates of



FIG. 8: The upper-left half-plane. Open stars are roots of the spanning trees on the white sublattice.

white sublattice on the cut are half-integers (see Fig.8). To avoid complication of notations, we shift the origin of the white sublattice to the point $(0,0)'$ shown in Fig.8.

After the transformation, the spanning trees on the sublattices belong to the upper-left half-space with respect to the diagonal cut. Also we can put the roots of the white sublattice at the line parallel to the cut and crossing the point $(0,0)'$ because all sites of this line can be connected with the root of the white spanning tree on the white sublattice.

III. LAPLACIAN AND GENERALIZED KIRCHHOFF THEOREM

Consider the graph $\mathcal{G} = (V, E)$ with vertex set V and set of bonds E. The vertices are sites of the square lattice and an additional point which is the root "$\star$": $V \equiv \{s_{x,y}, (x,y) \in \mathbb{Z}^2, |x| \leq M, |y| \leq N\} \cup \{\star\}$. The bonds of E connect neighboring sites of the lattice. We set the condition that all right and bottom boundary vertices $\{s_{x,+N}\}$ and $\{s_{-M,y}\}$ are connected with the root $\star$. The graph $\mathcal{G}$ represents the finite square lattice of size $(2N+1) \times (2M+1)$. We consider also the left-upper half plane $V_+ = \{s_{x,y}, x \leq y\}$ with open boundary conditions at the diagonal sites $V_0 = \{s_{x,y}, x = y\}$.

We construct the spanning tree by using the arrow representation. To this end, we attach to each vertex $i \in V \backslash \{\star\}$ an arrow directed along one of bonds $(i, i') \in E$ incident to it.

Each arrow defines a directed bond $(i \to i')$ and each configuration of arrows defines a spanning directed graph with set of bonds $\{(i \to i') : i \in V\backslash\{\star\},\ i' \in V,\ (i,i') \in E\}$. A sequence of directed bonds $(i_1, i_2),\ (i_2, i_3),\ (i_3, i_4), \ldots,\ (i_{m-1}, i_m)$ is called the path of length m from the site i_1 to the site i_m. This path forms a loop if $i_m = i_1$. No arrows are attached to the root $\star$, so that it is a sink of directed paths. Spanning tree is a spanning digraph without any loops. Our aim is to construct spanning trees, with a prescribed configuration of paths between fixed vertices. These configurations will be investigated by means of the determinant expansion of the Laplace matrix [11–13].

Let the vertices of the set V, be labeled in arbitrary order from 1 to $n = |V\backslash\{\star\}| = (2M+1)(2N+1)$. Then Laplacian Δ of size $n \times n$ has the elements:

$$\Delta_{ij} = \begin{cases} z_i & \text{if } i = j, \\ -1 & \text{if } i, j \text{ are nearest neighbors,} \\ 0 & \text{otherwise,} \end{cases} \tag{3.1}$$

where z_i is the degree of vertex $i \in V\backslash\{\star\}$. The determinant of Δ is a sum over all permutations σ of the set $\{1, 2, \ldots, n\}$:

$$\det \Delta = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \Delta_{1,\sigma(1)} \Delta_{2,\sigma(2)} \ldots \Delta_{n,\sigma(n)}\ , \tag{3.2}$$

where S_n is the symmetric group, $\operatorname{sgn}(\sigma) = \pm 1$ is the signature of permutation σ. In general, each permutation $\sigma \in S_n$ can be factorized into a composition of disjoint cyclic permutations, $c_1, c_2, \ldots c_k$. This representation partitions the set of vertices $V\backslash\{\star\}$ into non-empty disjoint subsets which are orbits $\mathcal{O}_i = \{v_{i,1}, v_{i,2} \ldots, v_{i,l_i}\} \subset V$ of the corresponding cycles c_i, $i = 1, \ldots, k$, at that $\cup_{i=1}^{k} \mathcal{O}_i = V\backslash\{\star\}$ and $\sum_{i=1}^{k} l_i = n$, where l_i is the length of cycle c_i. Orbits consisting of just one element are the fixed points $S_{fp}(\sigma)$ of the permutation. A cycle c_i of length $|c_i| = l_i \geq 2$ is called a proper cycle. The proper cycles are of even length only, hence, the number of proper cycles p defines the signature of the permutation σ, that is $\operatorname{sgn}(\sigma) = (-1)^p$. Thus (3.2) can be written as follows:

$$\det \Delta = \prod_{i=1}^{n} z_i + \sum_{p=1}^{[n/2]} (-1)^p \sum_{\sigma = \{c_1, \ldots, c_p\}} \prod_{i=1}^{p} \Delta_{v_i, c_i(v_i)} \Delta_{c_i(v_i), c_i^2(v_i)} \ldots \Delta_{c_i^{l_i-1}(v_i), v_i} \prod_{j \in S_{fp}(\sigma)} z_j\ , \tag{3.3}$$

where c_i^k is the k-fold composition of the cyclic permutation c_i of even length l_i, $v_i \in \mathcal{O}_i(\sigma)$, so that $c_i^{k-1}(v_i) \neq c_i^k(v_i)$ and $c_i^{l_i}(v_i) = v_i$. The term $\prod_{i=1}^{n} z_i$ equals to the number of all spanning digraphs having the root $\star$. Each of other terms on the right-hand side of (3.3) having a non-zero set of fixed points equals up to a sign to $\prod_{j\in S_{fp}(\sigma)} z_j$, because all non-diagonal elements equal to -1. That product represents the number of distinct spanning digraphs which have in common the specified cycles $c_1,\dots,c_p$, and differ in the oriented edges outgoing from vertices $j\in S_{fp}(\sigma)$. The expansion (3.3) can be interpreted in form of the inclusion-exclusion principle [20]. Let $c_1,c_2,\dots,c_m$ be the list of all possible proper cycles. We define A_i, $i=1,2,\dots,m$ as the set of all spanning digraphs containing the particular cycle c_i and A_0 is the set of all spanning digraphs. Let A_{ST} be the set of spanning trees i.e. the set of spanning digraphs containing no cycles. Then we can write down (3.3) in the form:

$$\det\Delta = |A_{ST}| = |A_0| - \sum_{i=1}^{m}|A_i| + \sum_{1\le i<j\le m}|A_i\cap A_j| + \ldots + (-1)^m|A_1\cap\ldots\cap A_m|\,, \quad (3.4)$$

where $|A|$ is cardinality of the set A. (3.4) is the Kirchhoff theorem for the number of spanning trees of a given graph [20].

Now we modify the Laplace matrix changing some non-diagonal elements:

$$\Delta'_{ij} = \begin{cases} z_i & \text{if } i=j,\\ -1 & \text{if } i,j \text{ are nearest neighbors},\\ -\varepsilon & \text{if } (i,j)\in\mathcal{B}\equiv\{(i_1,j_1),\ (i_2,j_2),\ldots(i_n,j_n)\},\\ 0 & \text{otherwise,}\end{cases} \quad (3.5)$$

where $\mathcal{B}$ is the set of directed bonds that are not nearest neighbors on the square lattice. The determinant expansion (3.3) with $\Delta^{'}$ generalizes the Kirchhoff theorem [12]. The product $\Delta^{'}_{i_1,j_1}\ldots\Delta^{'}_{i_n,j_n}=(-\varepsilon)^n$ survives in the limit $\lim_{\varepsilon\to\infty}\det\Delta'/\varepsilon^n$. Permutations corresponding to these terms, contain cycles with directed bonds $\mathcal{B}$. If the configuration of bonds $\mathcal{B}$ excludes cycles containing more than one bond from the set $\mathcal{B}$, the expression $\lim_{\varepsilon\to\infty}\det\Delta'/(-\varepsilon)^n$ gives the number of configurations with following features: (i) each configuration is the $(n+1)$- component spanning graph; (ii) k-th component consists of the path from site j_k to site i_k, $1\le k\le n$, and branches of the spanning tree attached to these paths; (iii) $(n+1)$-th component is the spanning tree containing the root $\star$. The ratio of numbers of such configurations and all one-component spanning trees is:

$$\lim_{\varepsilon\to\infty}\frac{\det\Delta'}{(-\varepsilon)^n\det\Delta} = \lim_{\varepsilon\to\infty}\frac{\det(I+\delta G)}{(-\varepsilon)^n}\,, \quad (3.6)$$

where $\delta = \Delta' - \Delta$ is the defect matrix and $G = \Delta^{-1} = G_{(p_1,q_1),(p_2,q_2)}$ is the Green function, which is defined in thermodynamical limit $M, N \to \infty$ on the plane as:

$$\begin{aligned} G_{(p_1,q_1),(p_2,q_2)} &= G_{p_2-p_1,q_2-q_1} = \\ &= G_{0,0} + \iint_{-\pi}^{\pi} \frac{\mathrm{d}\alpha \mathrm{d}\beta}{8\pi^2} \frac{e^{\mathrm{i}(p_1-p_2)\alpha+\mathrm{i}(q_1-q_2)\beta} - 1}{2 - \cos\alpha - \cos\beta} \,. \end{aligned} \tag{3.7}$$

If $\mathcal{B}$ includes m bonds $(i \to i')$ between nearest neighbors on the square lattice, the diagonal elements $\Delta'_{i,i}$ are changed from z_i to ε and non-diagonal ones $\Delta'_{i,i'}$ from -1 to $-\varepsilon$. In this case, the limit $\lim_{\varepsilon\to\infty} \det \Delta'/\varepsilon^m$ enumerates the spanning trees with an obligatory presence of bonds $(i \to i')$.

IV. CLUSTER PROBABILITIES

Upon above preparations, we are ready to calculate probabilities $P_a = P_{1a} + P_{2a} + P_{3a}$ and $P_b = P_{1b} + P_{2b} + P_{3b} - P_a$.

A. The probability P_a

The two-bond configuration corresponding to P_a is shown in Fig.9. We denote by B the non-zero submatrix of the defect matrix δ. For the case $1a$, matrix B is

$$B(\{1 \to 3, 3 \to 2\}) = \begin{pmatrix} \varepsilon & 0 & -\varepsilon \\ 0 & 0 & 0 \\ 0 & -\varepsilon & \varepsilon \end{pmatrix} . \tag{4.1}$$

The probability P_{1a} follows from (3.6)

$$P_{1a} = \lim_{\varepsilon\to\infty} \frac{\det(I + BG^{\mathrm{op}})}{(-\varepsilon)^2} \,, \tag{4.2}$$

where we changed δ by its non-zero finite submatrix B and the bulk Green function G by the Green function G^{op} for the open boundary conditions

$$G^{\mathrm{op}}_{(p_1,q_1),(p_2,q_2)} = G_{p_2-p_1,\, q_2-q_1} - G_{p_2-q_1,\, q_2-p_1} \,. \tag{4.3}$$

Using exact values (5.6) for the Green functions at short distances given in Appendix, we obtain

$$P_{1a} = \frac{16\pi - 16 - 3\pi^2}{6\pi^2} \,. \tag{4.4}$$



FIG. 9: Two-bond configuration.

For the case $3a$ we have

$$B(\{1 \to 3, 2 \to 3\}) = \begin{pmatrix} \varepsilon & 0 & -\varepsilon \\ 0 & \varepsilon & -\varepsilon \\ 0 & 0 & 0 \end{pmatrix} \tag{4.5}$$

and

$$P_{3a} = \frac{2\,(7\pi - 12 - \pi^2)}{\pi^2}\,. \tag{4.6}$$

The resulting probability P_a is

$$P_a = P_{3a} + 2P_{1a} = \frac{(4-\pi)(9\pi - 22)}{3\pi^2} = 0.181903\ldots \tag{4.7}$$

instead of $P_a = 1/4$ for the dense $O(1)$ loop model.

B. The probability P_b

The probability P_{1b} is the sum of two terms $P_{1b}(short)$ and $P_{1b}(long)$. The first of them is the probability that the self-avoiding path starting at site $(-1,0)$ of the original white sublattice stops at site $(1,0)$. The configuration of this path on the rotated sublattice is shown in Fig.10, where 1 on the original (non-rotated) lattice is the site $(-1,0)$ and 2 is the site $(1,0)$. Two bold arrows denote two possible connections of the site 2 with the roots. To generate the path from 1 to 2, we introduce the bridge $(2 \to 1)$ with the weight $-\varepsilon$ (see Fig.10) and tend ε to infinity. Then the defect matrix B consists of the single element



FIG. 10: The configuration corresponding to $P_{1b}(short)$ and its bridge representation.

$(2 \to 1)$ and the formula (3.6) is reduced to

$$P_{1b}(short) = 2 \lim_{\varepsilon \to \infty} \frac{\det(I + BG^{op})}{-\varepsilon} = 2(G_{1,1} - G_{2,1})\,. \tag{4.8}$$

Using the table (5.6) we get

$$P_{1b}(short) = \frac{2(\pi - 3)}{\pi}. \tag{4.9}$$

The probability $P_{1b}(long)$ involves the self-avoiding paths which start at the site $(-1,0)$, visit $(1,0)$ and then continue through the white sublattice up to the stop at a white site on the cut. This situation on the rotated sublattice is depicted in Fig.11, where points 1, 2, 3, 4 with the original coordinates are correspondingly $(-1,0)$, $(1,0)$, $(0,1)$, $(2,1)$. The path from the point 4 continues up to its end at a white site on the cut. We generate this path as



FIG. 11: The configuration corresponding to $P_{1b}(long)$ and its bridge representation.

a sum of two parts: one path from 1 to 2 via 3 and then the path from 4 to the ending point 5 (see Fig.11). Each of two parts appears due to auxiliary bridges (2 → 1) and (4 → 5). The orientations of the bridges ensure that a single path through two auxiliary bridges is forbidden. Then the formula (3.6) can be applied with the result

$$P_{1b}(long) = \frac{(\pi - 2)(16 - 5\pi)}{2\pi^2}\,, \tag{4.10}$$

so that

$$P_{1b} = P_{1b}(short) + P_{1b}(long) = \frac{14\pi - 32 - \pi^2}{2\pi^2}\,. \tag{4.11}$$

By the symmetry, we have $P_{1b} = P_{2b}$.

The crucial point of our calculations is the probability P_{3b}. The configuration of paths on the white sublattice corresponding to the case $3b$ gets after the transformation the form depicted in Fig.12, where the point 1 is the transformed position of the site $(-1, 0)$, the



FIG. 12: The configuration corresponding to case $3b$.

point 2 of the site $(1, 0)$ and the point 4 is the end of a path from the connection point s_0. In principle, it is possible to define a system of auxiliary bridges giving all three paths on Fig.12, however, in this case we would obtain a product of three Green functions which should be summed up over all positions of the point s_0. We choose here another way, avoiding these tremendous calculations.

First, we define the path from point 1 to point 4 inserting the auxiliary bond (4 → 1). Then, we fix an outgoing bond from point 2 in one of two possible directions (diagram I shown in Fig.13). The path from point 2 can join the path from point 1 to 4, forming thereby a configuration of the type 3b. Otherwise, the path from 2 can finish at one of the white sites on the boundary between points 2 and 4. The latter case, that is shown on the diagram II, Fig.14, must be excluded from all configurations in the diagram I. Thus, instead of summation over two-dimensional positions of the connection point s_0, we get two one-dimensional summations over positions of points 3 and 4, and reduce by one the number of Green functions involved into calculations.



FIG. 13: The diagram I representing P_I.

Consider now the Green function $G^{\rm op}_{(p,q),(k,k+1)}$ between an arbitrary site (p,q), $q>p$ on the left-upper half part of the square lattice and a site $(k,k+1)$ on the diagonal boundary. The summation over the whole boundary gives

$$\sum_{k=-\infty}^{+\infty} G^{\rm op}_{(p,q),(k,k+1)} = \sum_{k=-\infty}^{+\infty}\Big(G_{k-p,\,k+1-q} - G_{k-q,\,k+1-p}\Big) = \tag{4.12}$$

$$\frac{1}{8\pi^2}\iint_{-\pi}^{\pi}\sum_{k=-\infty}^{+\infty}\frac{e^{{\rm i}\,k(\alpha+\beta)}e^{{\rm i}\,\beta}\big(e^{-{\rm i}\,(p\alpha+q\beta)} - e^{-{\rm i}\,(q\alpha+p\beta)}\big)}{2-\cos\alpha-\cos\beta}\,{\rm d}\alpha\,{\rm d}\beta =$$

$$= \frac{1}{4\pi}\int_{-\pi}^{\pi}\frac{{\rm i}\,e^{-{\rm i}\,\alpha}\sin[(q-p)\alpha]}{1-\cos\alpha}\,{\rm d}\alpha = \frac{1}{4\pi}\int_{-\pi}^{\pi}\frac{\sin\alpha\sin[(q-p)\alpha]}{1-\cos\alpha}\,{\rm d}\alpha = \frac{1}{2}\,. \tag{4.13}$$



FIG. 14: The diagram II representing P_{II}.

With $p = 0$ and $q = 1$ we have

$$D_k \equiv G^{\rm op}_{(0,1),(k,k+1)} = G_{k,k} - G_{k-1,k+1} \ , \tag{4.14}$$

and

$$\sum_{k=-\infty}^{+\infty} D_k = \frac{1}{2} \ . \tag{4.15}$$

Due to the symmetry relation

$$D_{-k} = D_k \ . \tag{4.16}$$

we can find the sum

$$\sum_{k=1}^{+\infty} D_k = \frac{1}{4} - \frac{1}{2}(G_{0,0} - G_{1,1}) = \frac{1}{4} - \frac{1}{2\pi} \ . \tag{4.17}$$

Denote by R_k the probability of the diagram I for distance k between points 1 and 4. The calculation of the determinants with defect matrices B containing the auxiliary bond between points 1 and 4, and selected outgoing bonds from the point 2 gives

$$R_k = \frac{4(\pi-3)}{\pi} D_{k-1} + \frac{2(\pi-2)}{\pi} D_k \ . \tag{4.18}$$

The probability P_I is the sum

$$P_I = \sum_{k=2}^{\infty} R_k \ . \tag{4.19}$$

From the identity

$$\sum_{k=-\infty}^{+\infty} R_k = 2\sum_{k=2}^{+\infty} R_k + \frac{4(\pi-3)}{\pi} D_0 + \frac{2(\pi-2)}{\pi}\left(D_{-1}+D_0+D_1\right) \tag{4.20}$$

we have

$$P_I = \frac{6\,\pi - 8 - \pi^2}{2\,\pi^2}. \tag{4.21}$$

The probability P_{II} of the diagram II (see Fig.14) is a double sum of double bridges with distance s between points 1 and 4 and distance k between points 1 and 3. The corresponding B matrix reads

$$B = \begin{pmatrix} 0 & 0 & 0 & -\varepsilon \\ 0 & 0 & -\varepsilon & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \tag{4.22}$$

which gives the probability

$$P_{II} = 4 \lim_{\varepsilon\to\infty} \frac{\det(I + BG^{op})}{\varepsilon^2} = 4 \sum_{2\le k<s} \left(D_{k-1}D_s - D_k D_{s-1}\right) = \tag{4.23}$$

$$= 4 \sum_{2\le k\le s} D_s(D_{k-1} - D_k) + 4\sum_{k=2}^{\infty}\sum_{s=k}^{\infty} D_k(D_s - D_{s-1}) = \tag{4.24}$$

$$= 4\sum_{s=2}^{\infty} D_s(D_1 - D_s) - 4\sum_{k=2}^{\infty} D_k D_{k-1} = \tag{4.25}$$

$$= 4D_1 \sum_{s=2}^{\infty} D_s - 4\sum_{k=2}^{\infty} D_k(D_k + D_{k-1})\ , \tag{4.26}$$

or, in a more convenient form,

$$P_{II} = 4D_1 \sum_{k=1}^{+\infty} D_k - 4\sum_{k=1}^{+\infty} D_k(D_k + D_{k+1})\ . \tag{4.27}$$

One can show that

$$\sum_{k=-\infty}^{+\infty} D_k(D_k + D_{k+1}) = 2\sum_{k=1}^{+\infty} D_k(D_k + D_{k+1}) + D_0(D_0 + 2D_1) \tag{4.28}$$

so

$$P_{II} = 4D_1 \sum_{k=1}^{+\infty} D_k + 2D_0(D_0 + 2D_1) - 2\sum_{k=-\infty}^{+\infty} D_k(D_k + D_{k+1})\ . \tag{4.29}$$

Denote the latter sum by Q:

$$Q = \sum_{k=-\infty}^{+\infty} D_k(D_k + D_{k+1})\ . \tag{4.30}$$

Using the integral representation of the Green function and definition of D_k, after some manipulations we can show that

$$Q = \frac{1}{16\,\pi^3} \iiint_{-\pi}^{\pi} \frac{(\sin\alpha_1 - \sin\beta_1)(\sin\alpha_1 - \sin(2\alpha_1 + 2\alpha_2 + \beta_1))}{(2 - \cos\alpha_1 - \cos\beta_1)(2 - \cos\alpha_2 - \cos(\alpha_1 + \beta_1 + \alpha_2))} \, \mathrm{d}\alpha_1 \, \mathrm{d}\beta_1 \, \mathrm{d}\alpha_2 \, . \quad (4.31)$$

By the substitution

$$t_1 = \tan\frac{\alpha_1}{2}, \qquad t_2 = \tan\frac{\beta_1}{2}, \qquad t_3 = \tan\frac{\alpha_2}{2}, \quad (4.32)$$

$$\sin\alpha_1 = \frac{2t_1}{1+t_1^2}, \quad \sin\beta_1 = \frac{2t_2}{1+t_2^2}, \quad \sin\alpha_2 = \frac{2t_3}{1+t_3^2}, \quad (4.33)$$

$$\cos\alpha_1 = \frac{1-t_1^2}{1+t_1^2}, \quad \cos\beta_1 = \frac{1-t_2^2}{1+t_2^2}, \quad \cos\alpha_2 = \frac{1-t_3^2}{1+t_3^2}, \quad (4.34)$$

$$\mathrm{d}\alpha_1 \, \mathrm{d}\beta_1 \, \mathrm{d}\alpha_2 = \frac{2}{1+t_1^2}\frac{2}{1+t_2^2}\frac{2}{1+t_3^2} \, \mathrm{d}t_1 \, \mathrm{d}t_2 \, \mathrm{d}t_3 \quad (4.35)$$

and symmetrization by permutation $t_1 \rightleftarrows t_2$, we come to the triple integral of a rational function

$$Q = \frac{1}{\pi^3} \iiint_{-\infty}^{\infty} T_1(t_1, t_2, t_3) T_2(t_1, t_2, t_3) \, \mathrm{d}t_1 \, \mathrm{d}t_2 \, \mathrm{d}t_3 \, , \quad (4.36)$$

where

$$T_1(t_1, t_2, t_3) = \frac{(t_1 - t_2)^2 (t_1 t_2 - 1)(t_1 t_2 + t_1 t_3 + t_2 t_3 - 1)(t_1 t_2 t_3 - t_1 - t_2 - t_3)}{(1+t_1^2)^2 (1+t_2^2)^2 (1+t_3^2)^2 (t_1^2 + t_2^2 + 2t_1^2 t_2^2)} \, , \quad (4.37)$$

$$T_2(t_1, t_2, t_3) = \frac{2t_1 t_2 t_3 - 2t_3 + (t_1 + t_2)(t_3^2 - 1)}{(t_1 + t_2 + t_3)^2 - 2t_1 t_2 t_3 (t_1 + t_2 + t_3) + 2t_1^2 t_2^2 t_3^2 + (1 + t_1^2 + t_2^2) t_3^2} \, . \quad (4.38)$$

After integration over t_3 and several manipulations we have

$$Q = \frac{2}{\pi^2} \int_0^{+\infty} \int_{-t_1}^{+t_1} \frac{(t_1 - t_2)^2 \left(1 + t_1^2 + t_2^2 + t_1^2 t_2^2 - (t_1 + t_2)\sqrt{(1+t_1^2)(1+t_2^2)}\right)}{(1+t_1^2)^2 (1+t_2^2)^2 (t_1^2 + t_2^2 + 2t_1^2 t_2^2)} \, \mathrm{d}t_2 \, \mathrm{d}t_1 \, . \quad (4.39)$$

The integration over t_2 gives

$$Q = \frac{1}{\pi^2} \int_0^{+\infty} \left(\frac{8t_1^3 \arctan\left(\sqrt{1+2t_1^2}\right)}{(1+t_1^2)^2 \sqrt{1+2t_1^2}} + \frac{4(1-t_1^2)\arctan(t_1)}{(1+t_1^2)^2} - \frac{2(\pi - 2)t_1^2}{(1+t_1^2)^2} \right) \mathrm{d}t_1 \, , \quad (4.40)$$

from which we obtain

$$Q = \frac{4 - \pi}{2\pi} \, . \quad (4.41)$$

As a result, we have

$$P_{II} = \frac{2\pi^2 - 5\pi - 4}{\pi^2} \, . \quad (4.42)$$

and

$$P_{3b} = 2 \times P_I - 2 \times P_{II} = \frac{16 - 5\pi}{\pi}\,. \tag{4.43}$$

The final result for P_b is

$$P_b = P_{1b} + P_{2b} + P_{3b} - P_a = \frac{32\pi - 8 - 9\pi^2}{3\pi^2}\,. \tag{4.44}$$

The numerical value of this probability is $P_b = 1/8 + 0.0001156...$ that is remarkably close to the conjectured value $P_b = 1/8$ for the dense $O(1)$ loop model.

The calculation of the minimal link probability for $n = 0$ is a first step in a program of investigations of more general link correlation functions. Being realized, this program would help to establish a correspondence between lattice calculations and operators of the logarithmic conformal field theory (LCFT) with the central charge $c = -2$. Then, the operators of the LCFT corresponding to link variables in the dense $O(1)$ loop model with the central charge $c = 0$ can be constructed by an analogy with the $O(0)$ model.

V. APPENDIX: GREEN FUNCTIONS

The explicit form of the translationally invariant Green function on the plane is

$$G_{\vec{r}_1,\vec{r}_2} \equiv G(\vec{r}_2 - \vec{r}_1) \equiv G_{0,0} + g_{p,q}, \quad \vec{r}_2 - \vec{r}_1 \equiv \vec{r} \equiv (p,q) \tag{5.1}$$

with $G_{0,0}$ an irrelevant infinite constant. The (finite) numbers $g_{p,q}$ are given explicitly by

$$g_{p,q} = \frac{1}{8\pi^2} \iint_{-\pi}^{\pi} \frac{e^{\mathrm{i}\,p\,\alpha + \mathrm{i}\,q\,\beta} - 1}{2 - \cos\alpha - \cos\beta}\, \mathrm{d}\alpha\, \mathrm{d}\beta\,. \tag{5.2}$$

Let us mention symmetry properties of this function:

$$g_{p,q} = g_{q,p} = g_{-p,q} = g_{p,-q}\,. \tag{5.3}$$

After the integration over α, it can be expressed in a more convenient form for actual calculations,

$$g_{p,q} = \frac{1}{4\pi} \int_{-\pi}^{\pi} \frac{t^p\, e^{\mathrm{i}\,q\,\beta} - 1}{\sqrt{y^2 - 1}}\, \mathrm{d}\beta\,, \tag{5.4}$$

where $t = y - \sqrt{y^2 - 1}$, $y = 2 - \cos\beta$ (see [13]). For $r^2 = p^2 + q^2 \gg 1$ it has a behavior [18]

$$g_{p,q} = -\frac{1}{2\pi}\left(\log r + \gamma + \frac{3}{2}\log 2\right) + \frac{\cos(4\,\varphi)}{24\,\pi\, r^2} + \frac{18\cos(4\,\varphi) + 25\cos(8\,\varphi)}{480\,\pi\, r^4} + \ldots\,, \tag{5.5}$$

where $(p,q) = (r\cos\varphi, r\sin\varphi)$, and with $\gamma = 0.57721\ldots$ the Euler constant. Let us also write down some values of the Green functions for particular points (p,q), which are needed for calculations:

$$\begin{array}{lll} g_{0,1} = -\frac{1}{4} & g_{0,2} = -1 + \frac{2}{\pi} & g_{0,3} = -\frac{17}{4} + \frac{12}{\pi} \\ g_{1,1} = -\frac{1}{\pi} & g_{1,2} = \frac{1}{4} - \frac{2}{\pi} & g_{1,3} = 2 - \frac{23}{3\pi} \\ g_{2,2} = -\frac{4}{3\pi} & g_{2,3} = -\frac{1}{4} - \frac{2}{3\pi} & g_{3,3} = -\frac{23}{15\pi}\,. \end{array} \tag{5.6}$$

Acknowledgments

This work was supported by the RFBR grants No. 12-01-00242a, 12-02-91333a, the Heisenberg-Landau program, the DFG grant RI 317/16-1 and the grant of NRU Higher School of Economics, Academic Fund program No.12-09-0051. V.B.P. thanks Vladimir Rittenberg for useful discussions. V.S.P. thanks Philippe Ruelle for hospitality in UCL (Louvain-la-Neuve, Belgium) and critical remarks.

---

[1] H.W.J. Blöte and B. Nienhuis, J. Phys. A **22**, no. 9, 1415 (1989).

[2] Yu.G. Stroganov, J.Phys. A **34**, no.13, L179 (2001).

[3] A.V. Razumov and Yu.G. Stroganov, J. Phys. A **34**, no.14, 3185 (2001).

[4] A.V. Razumov and Yu.G. Stroganov, Theor. Math. Phys. **138**, 333 (2004).

[5] M.T. Batchelor, J. de Gier and B. Nienhuis, J.Phys. A **34**, no.19, L265 (2001).

[6] J. de Gier, B. Nienhuis, P.A. Pearce and V. Rittenberg, Phys. Rev. E **67**, 016101 (2003).

[7] S. Mitra, B. Nienhuis, J. de Gier and M.T. Batchelor, J. Stat. Mech.:Theor.Exp. P09010 (2004).

[8] G. Kuperberg, Ann. of Math. (2) **156**, no.3, 835 (2002).

[9] P.A. Pearce and J. Rasmussen, J. Stat. Mech.:Theor.Exp. (2007) P02015.

[10] G. Kirchhoff, Ann. Phys. Chem., **72**, 497 (1847).

[11] D. Dhar, Phys. Rev. Lett. **64**, 1613 (1990).

[12] V.B. Priezzhev, J. Stat. Phys. **74**, 955 (1994).

[13] G. Piroux and P. Ruelle, Phys. Lett. B **607** (2005) 188;
M. Jeng, G. Piroux and P. Ruelle, J. Stat. Mech. (2006) P10015.

[14] V.S. Poghosyan, S.Y. Grigorev, V.B. Priezzhev and P. Ruelle, J. Stat. Mech. (2010) P07025.

[15] S. Caracciolo and A. Sportiello, J. Stat. Mech. (2012) P09013

[16] L. Levine and Y. Peres, arXiv:1106.2226, published in Random Structures and Algorithms.

[17] V.S. Poghosyan, V.B. Priezzhev and P. Ruelle, J. Stat. Mech.:Theor.Exp. (2011) P10004.

[18] S.Y. Grigorev, V.S. Poghosyan and V.B. Priezzhev, J. Stat. Mech.:Theor.Exp. (2009) P09008.

[19] A. Gorsky, S. Nechaev, V.S. Poghosyan and V.B. Priezzhev arXiv:1206.3147.

[20] V.B. Priezzhev, Sov.Phys.Usp. **28**, (1125) 1985.